### Particulate Inorganic Carbon

William Balch
Bigelow Laboratory for Ocean
Sciences
E. Boothbay, ME 04544


bbalch@bigelow.org





### Acknowledgments

- Bruce Bowler, Dave Drapeau and Laura Lubelczyk, (Bigelow Laboratory for Ocean Sciences)
- Captains and crew of: RRS Discovery, R/V Melville, R/V Revelle, R/V Ron Brown, RRS James Clark Ross, RRS James Cook, R/V Thompson; R/V Tangaroa; M/S Scotia Prince; M/S The CAT
- Norman Kuring- NASA Goddard
- Goddard Ocean Color Group
- Giovanni and Ocean Data View
- NASA support



# PIC algorithm is typically focused on coccolithophores- ubiquitous, single-cell calcifying plants




There are other PIC particles in the sea but coccoliths and coccolithophores have the highest scattering per unit PIC ("backscattering cross section")

#### **Outline**

- PIC algorithm cross platform comparisons
- PIC algorithm refinement


Science applications





### Cross platform algorithm variability

- SeaWiFS
- MODIS Terra
- MODIS Aqua

Merged Two-Band/Three band algorithm for PIC applied to SeaWiFS and **MODIS-Aqua** data (Collection 6)...Similar patterns









#### PIC Algorithm Performance: Collection 6

Scatter Plot (Time Averaged)
Time: Jul2002—Dec2012 Area: (905—90N, 180W—180E)




MODIS Aqua PIC (9km; mol m<sup>-3</sup>)


#### MODIS Aqua PIC, on average, < SeaWiFS



SeaWiFS Aqua PIC (9km; mol m<sup>-3</sup>)



We've used MISR to measure coccolith backscattering... using the Blue bands, integrated -70° to 70°, subtracting the NIR band (quasi atmosphere correction), then a dark pixel assumption to further correct.



### PIC algorithm refinement

- The backscattering that we associated with chlorophyll has been adjusted for PIC.
- The backscattering cross section of PIC (b<sub>b</sub>\*; m<sup>2</sup> mol<sup>-1</sup>) has been adjusted (based on cruise work).
   Variable b<sub>b</sub>\* algorithm.
- New K<sub>490</sub> formulation in PIC algorithm (formerly Baker and Smith 1982; now Morel and Martorena, 2001)
- F<sub>o</sub> has been updated to what is used in SeaDAS

#### Optical Modeling of the Upper Ocean in Relation to Its Biogenous Matter Content (Case I Waters)







Fig. 7a. A log-log plot of the difference  $K(450) - K_w(450)$  as a function of the pigment concentration for case I waters only. The straight lines correspond to linear fits obtained by regression, whereas the curve comes from *Baker and Smith* [1982]. The case II waters, excluded from regression, are not shown.

# Many major cruises to study coccolithophores- both northern and southern hemisphere; sampled over >212,000 km...





#### b<sub>b</sub>\* highly variable; some latitudinal variability





### LUT changes








Satlantic Micro-SAS radiometers collect along-track data for Lt, Lsky, and Ed. We designed a mount which compensated real-time for changes in sun's azimuth relative to ship heading.


Radiometers mounted on the RRS Discovery

Balch; Bigelow Lab

### Algorithm improvement (n=~800)




### New algorithm changes (LUT & bb\*)




Particulate Inorganic Carbon Mean [mol m^-3]

### Relating Surface to integrated PIC...




### Science applications


**Using MODIS** SST, Aquarius SSS to define global surface density field. High PIC sits at the strong gradients of **Sub Antarctic Frontal** boundary




Global Surface Density (kg m-3)




Plot surface PIC concentration in temperature/salinity space PIC (mol m<sup>-3</sup>




### PIC Global Time Series (MODIS-Aqua) Mission record




# Global Surface PIC (MODIS-Aqua) Mission record




# Global Int. PIC (MODIS-Aqua) Mission record



## PIC Global Time Series (MODIS-Aqua) Annual coccolithophore phenology



## PIC Global Time Series (MODIS-Aqua) Annual coccolithophore phenology



## Implications of these results to PIC turnover...

- Average global PIC =~5[+/-1] x 10<sup>11</sup> moles PIC
   =6 x 10<sup>12</sup> g PIC
- Global calcification = 1.6[+/-0.3] Pg PIC y<sup>-1</sup>
   Balch et al., 2007
- $(6 \times 10^{12} \text{ g PIC})/(1.6 \times 10^{15} \text{ g PIC y}^{-1}) = 0.003 \text{ y}$ =~1.1d turnover
- What is happening to it? It sinks or dissolves
- No indication of a trend in global PIC (e.g. negative trend associated with OA)

### Summary- PIC algorithm

- Cross platform comparisons of PIC are within 3-13% depending on the platform
- MISR can be used to detect coccolithophores
- New improved PIC algorithm has <u>accuracy of</u> +/-0.2-0.3mmol PIC m<sup>-3</sup>= 2-3μg PIC L<sup>-1</sup>
- PIC is not homogenously distributed but shows subsurface maxima in oligotrophic waters, and surface maxima in blooms; use this to improve vertical integrations based on surface PIC estimates

### Summary- Science Issues

- Highest [PIC] seen globally at frontal boundaries between surface seawater densities ( $\sigma_{\theta}$ ) 26-27 (SSS 34-36 and SST 2-15°C)
- Greatest global PIC standing stock in January and lowest in March
- Global phenology of coccolithophore blooms shows greatest average [PIC] associated with austral summer, boreal summer lower, lowest at the equinoxes

### Future tasks at hand; Improve PIC algorithm performance

- Examine using reflectancedifferencing algorithm (Hu et al., 2012) for determining b<sub>b chl</sub>
- Improve predictions of global variability of b<sub>b</sub>\*
- Partition backscattering into b<sub>b POC</sub>, b<sub>b</sub>
   PIC and b<sub>b BSi</sub>


### Future: MODIS Science

- Examine coccolithophore bloom at subkilometer pixel resolution using MISR and high-resolution land bands. What can we learn about processes driving bloom at these scales?
- Examine global coccolithophore
   phenology in different portions of world ocean (different biogeochemical provinces)
- Frontal impacts on <u>coccolithophore growth</u> (using Aquarius salinity; MODIS SST)

### Future: MODIS Science


- Impact of wind speed on coccolithophore standing stocks (AMSR-E or Aquarius winds)
- Examine <u>PIC Turnover</u> in different regions of world <u>as foci for ballasted sinking POC</u>
- Are there <u>long-term negative trends of PIC</u> <u>associated with Ocean Acidification</u>, especially in polar provinces where OA will have biggest effects?

### Thank you!




# While satellites generally agree with collection 6, shipboard validation data show underestimates...the issue is b<sub>h</sub>\*

GB II and AMT22 PIC vs bb' (within 0.5 hr)




Sampled within ½ h of overpass


# Use shipboard radiance data to predict PIC bb...




### Global Surf. PIC Anomaly (MODIS-Aqua)



### Global Int. PIC Anomaly (MODIS-Aqua)




#### MODIS Terra PIC, on average, ~6% > MODIS Aqua



MODIS Aqua PIC (9km; mol m<sup>-3</sup>)

#### MODIS Terra, on average, within ~3% of SeaWiFS



SeaWiFS PIC (9km; mol m<sup>-3</sup>)

### Ocean area viewed varies annualy

